

FOR OFFICIAL USE.

M.O. 177 (1917).

EXAMINED BY  
AUTHORITY FOR ISSUE  
ISSUED ON

250 Recd 2/8/17 - J.H.B.T.

# THE RELATION

BETWEEN

## PRESSURE, TEMPERATURE, AND AIR CIRCULATION OVER THE SOUTH ATLANTIC OCEAN.

NOTES WITH REFERENCE TO A SET OF MONTHLY WIND CHARTS OF THE SOUTH ATLANTIC OCEAN WHICH WERE PREPARED IN THE METEOROLOGICAL OFFICE AND WERE ORIGINALLY PUBLISHED BY THE HYDROGRAPHIC DEPARTMENT OF THE ADMIRALTY, IN JANUARY, 1904.

SECOND EDITION.

BY

M. W. CAMPBELL HEPWORTH, C.B., R.D.,  
*Captain R.N.R., Marine Superintendent.*

Published by the Authority of the Meteorological Committee.



LONDON:

PUBLISHED BY HIS MAJESTY'S STATIONERY OFFICE.

To be purchased through any Bookseller or directly from  
H.M. STATIONERY OFFICE at the following addresses:  
IMPERIAL HOUSE, KINGSWAY, LONDON, W.C.2, and  
28, ABINGDON STREET, LONDON, S.W.1;  
37, PETER STREET, MANCHESTER; 1, ST. ANDREW'S CRESCENT, CARDIFF;  
23, FORTH STREET, EDINBURGH;  
or from E. PONSONBY, LTD., 116, GRAFTON STREET, DUBLIN;  
or from the Agencies in the British Colonies and Dependencies,  
the United States of America and other Foreign Countries of  
T. FISHER UNWIN, LTD., LONDON, W.C.

1917.

Price 1s. Net.

FG5  
Stock 8



## PREFACE.

The discussion of the meteorological observations for the South Atlantic and the South American Coastal Regions of the Pacific occupied the attention of the Marine Department of the Meteorological Office from 1898, when the corresponding work upon the Southern Ocean was completed, until 1904. The total number of observations extracted from the Log Books of the Royal Navy, and the Meteorological Logs contributed by the Captains and Officers of the Mercantile Marine whose ships crossed the region referred to, amounted to 946,000. They have been grouped according to five degree squares of latitude and longitude, or in special cases according to one degree squares, and the mean results for Wind, Air Pressure, and Air Temperature were embodied in a series of monthly charts prepared under Captain Hepworth's superintendence in accordance with the direction of the Meteorological Council and, in particular, of the late Hydrographer of the Navy, Rear-Admiral Sir W. J. L. Wharton, F.R.S., who was ex officio a member of the Council.

Upon the completion of the Charts the Hydrographer undertook the issue of them as a publication of the Hydrographic Office. The information contained in the Charts will, it is hoped, be found to be of considerable interest and importance to mariners. At the request of the Council, Captain Hepworth undertook the preparation of some notes giving the general results which follow from an examination of the Charts. They formed the subject matter of a paper read before Section A. of the British Association at Cambridge in 1904, and are now issued by the authority of the Meteorological Committee.

Charts for the Coastal Regions of South America were issued first in 1902, and the issue of the completed work for the South Atlantic followed in 1904.

By way of illustration of some of the general features of the average meteorological conditions of the South Atlantic, some additional Charts on small scale have been prepared. They show the variations in the course of the year of the position and intensity of the areas of high pressure in the regions under consideration, and the position and extent of the equatorial region

(11354-12.) Wt. 7107-874. 350. 7/17. D & S. G. 3.



3 8078 0001 5574 9



of doldrums in relation thereto, and the average distribution of gales and of fog. Captain Hepworth's notes call attention to the points which are illustrated by these additional Charts, and also to the effect of the air circulation upon the distribution of temperature, which is found to diverge considerably from the parallels of latitude, even over an area of such uniform geographical character as the South Atlantic.

For the facts upon which the notes depend, the Wind Charts themselves should be referred to, but it is thought that sailors and others interested in marine meteorology may wish to have the general conclusions in a separate form.

W. N. SHAW,  
Director.

Meteorological Office,  
63, Victoria Street, S.W.  
August, 1905.

NOTE.—27 March, 1917.—Not much has been added to our knowledge of the South Atlantic Ocean since the M.O. Charts were published in 1904, and the call for a new edition of Captain Hepworth's notes can therefore be met with some slight alterations of the text and the conversion of the pressure-readings in order to correct them for latitude in accordance with international convention, and to bring them into accordance with the practice of the Meteorological Office, adopted also since January, 1917, by the Bureau Central Météorologique of Paris, of expressing pressure in millibars.

NAPIER SHAW.

Meteorological Office,  
South Kensington, S.W.7.

## THE RELATION

BETWEEN

### PRESSURE, TEMPERATURE, AND AIR CIRCULATION OVER THE SOUTH ATLANTIC OCEAN.

#### WIND AND BAROMETER.

The South Atlantic offers an excellent field for the study of air circulation on a comprehensive scale and under normal conditions. An ocean covering an extensive area, connected north and south with great oceans and completely open to the south, it is at the same time free from the disturbing influences of island groups, and is bounded east and west by continents having coastlines that are, for the most part, exempt from large irregularities of outline. The atmospheric circulation over the South Atlantic may therefore be regarded as one vast wind system, its air currents undisturbed over the northern half by the occurrence of aerial eddies for the most part; and over the southern half, although at times interrupted, yet not effaced by them.

The relation between Pressure, Temperature, and Air Circulation, over this area, also over the eastern margin of the South Pacific Ocean, is well shown by a set of monthly wind charts prepared in the Meteorological Office and published by the Admiralty, entitled *Wind Charts for the South Atlantic Ocean*. These charts embody the general results of a very large number of observations of wind and simultaneous readings of the barometer and thermometer.

The anticyclone, or area of high bar. metric pressure, is the great feature of the system. Round this central high pressure a general circulation of air is indicated.

This circulation of air is assisted on the eastern side of the ocean by the relatively low pressure over South Africa; on the northern side by the low pressure over the equatorial regions of the Atlantic; on the western side by the relatively low pressure over South America; and on the southern by the lower pressure of higher latitudes. The direction of the wind about the core of high pressure, stated generally, is Southerly to South-easterly between it and the African coast; South-easterly and Easterly, towards the equator; Easterly to Northerly on its western side; and North-westerly to South-westerly on its southern. Over the eastern and northern segments of the system the flow of the air current is steady; over the western and southern, particularly over the former, the circulation is less persistent. Southward of 30° S. lat., and even farther to the northward, on the western side of the South Atlantic, the apparent normal circulation of the surface





wind, in harmony with the course of the average isobars, is largely masked by the effect of low pressure systems travelling eastward or south-eastward.

These low pressure systems occur more frequently from May to October, inclusive, than in other months, and are most frequent in July and August. South of  $40^{\circ}$  S. lat., the westerly winds, although temporarily interrupted by depressions on their approach, are intensified by them on their retreat. These depressions generally appear on the ocean between the 20th and 40th parallels, and the track of those approaching the margin of the relatively high pressure to the eastward is frequently turned south-eastward. The cool northward-flowing air current, on the eastern side of the ocean, reinforces the area of high pressure which acts as a barrier to eastward moving depressions; the warm southward-flowing air current, on the western side, induces conditions favourable to the formation or advance of disturbances in that region. Thus the most striking characteristic of the distribution of winds, shown on the charts referred to, is the steady south-easterly current on the eastern and northern sides of the tropical high pressure, and the varying winds on the western side. The varying character of the winds on the western side must be attributed to the passage of low pressure areas mentioned above across the region, or to the south of it. Observations in these latitudes are too few to enable us to identify the character or trace the track of these depressions over the ocean, but observations over the adjacent land area of South America occasionally show low pressure areas which pass away over the ocean not far from the localities referred to.

Four charts showing isobars over South America, *see* Plates I. and II., taken from the Daily Weather Reports issued by the Argentine Meteorological Office, may serve to illustrate the relation of the moving depressions to the general distribution of pressure over the South Atlantic. The dates given on the charts indicate the days on which the low pressure systems were observed; the dotted lines over the ocean show the distribution of mean pressure for the month. Areas of low pressure are shown, from time to time, by the Daily Weather Charts of Argentina, to reach the sea on a parallel as far north as  $20^{\circ}$  S. lat.; but observations are sparse between the 20th and 30th parallels, and no daily synoptic charts are published for Brazil.

Log books in the possession of the Meteorological Office, however, record numerous instances of strong winds associated with low pressure systems experienced in the vicinity of the Brazilian coast, between  $20^{\circ}$  S. and  $30^{\circ}$  S.,  $40^{\circ}$  W. and  $50^{\circ}$  W., many of which attained gale force. In the years 1855 to 1899 no less than 184 gales were recorded in these Registers, and in all of these instances the force of the wind reached, or exceeded, 8 of Beaufort scale, and the strongest gale recorded attained force 11 of that scale. The gale forces cannot be assigned to winds from any particular direction; they come from every quarter. The passage eastward of a cyclonic depression over the River Plate, or its neighbourhood, appears to occur with some regularity at about the end of August.



Illustrating the passage of low pressure systems over the Ocean. The Isobars east of the 40th meridian refer to normal pressure distributions.



On the eastern margins of the South Atlantic, and South Pacific north of  $35^{\circ}$  S. lat., and in a measure also on the western margin of the former, north of the same parallel, when the conditions of the atmosphere are stable, there appears a marked inclination for the wind to follow the course of the littoral. This probably is connected with the tendency of barometric pressure over the land to conform to the contour of the coast-line, resulting from a slight temperature and consequent pressure gradient, between land and sea.

Throughout the 12 months the shape of the isobar bounding the area of highest pressure, 1020 mb. (30.12 ins.) or 1023 mb. (30.21 ins.), is approximately that of an ellipse. The major axis, however, does not always make the same angle with the meridian; and the axes vary, not only in position, but also in length, from month to month.


Over the South Pacific throughout the 12 months there is an area of high barometer readings, the South Pacific anticyclone, which is elliptical in shape, and is indicated by an isobar of 1020 mb. (30.12 ins.) except in August, when the highest value shown is 1023 mb. (30.21 ins.).

Small charts showing the intensity and position of the region of high pressure over the South Atlantic and eastern margin of the Pacific in each month, are given on Plates III. to VI., and, as having a close connexion with the changes shown on these charts, the mean latitude of the belt of doldrums in the Atlantic is also indicated.

The general features of the distribution of pressure, wind, and temperature, at the various seasons of the year, are as follow:—

*December to February inclusive.*—Over the South Atlantic eastward from the area of highest pressure to the African coast; northward and north-westward to the equator, and coast of South America, as well as over its eastern and northern segments, a steady trade wind is found mainly from some point between S.E. and E., but the wind becomes a Southerly or even South-south-westerly on the African side, and North-easterly or even Northerly near South America where the trend of the coast-line is the dominant factor. To the westward of the area of highest pressure, stated broadly, the direction of the prevailing wind following the course of the isobars, is North-easterly, although depressions passing eastward and south-eastward partly mask the circulation conforming to the average distribution of pressure, and near the South American coast the influence of the land upon the direction of the wind can be clearly traced. South of the  $35^{\text{th}}$  parallel, which is about the latitude of Monte Video, the prevailing winds have some westing in them, and a good deal of nothing, except where they appear to be influenced by the land west of the  $40^{\text{th}}$  meridian between the  $35^{\text{th}}$  and  $40^{\text{th}}$  parallels.

South of the  $40^{\text{th}}$  parallel Westerly winds largely predominate in all months of the year; and the relation between the slope of the gradient and the strength of the wind is well shown by the percentage of gales for the regions where the isobars close up.



Illustrating the passage of low pressure systems over the Ocean.  
The Isobars east of the  $40^{\text{th}}$  meridian refer to normal pressure distributions.



Between the 40th and 55th parallels the pressure gradient is almost always towards the south, and Easterly winds are correspondingly rare.


Over the eastern margin of the South Pacific from 30° S. northward, the trade winds blow steadily; they are chiefly from a South-easterly direction, but near the South American Continent the prevailing winds flow parallel to the outline of the coast. Southward of 35° S., in accordance with the trend of the isobars, the air current is South-westerly, where not deflected by proximity to land; southward of 40° S. a Westerly air current predominates.

*March to June inclusive.*—Over the South Atlantic to the eastward of the high pressure area, as well as over its eastern and northern margins; and from 20° S. northward to the equator, the air current is mainly from some point between S. and E., except where deflected by land influence. Westward of the 20th meridian between 20° S. and 35° S. it has a good deal of northing and easting, but is not constant; and this area probably is visited by many small disturbances and is subject to varying influences. South of 35° S. westerly winds predominate.

Over the south-eastern margin of the South Pacific, the area over which the south-east trade winds prevail, is gradually reduced, so that in June they are not constant south of the 20th parallel. After March easterly winds become less and less frequent south of the 25th parallel; south of the 40th the westerly winds are well established.

*July to September inclusive.*—Over the South Atlantic there is a general resemblance between the highest pressure areas of the first and this the third period; but in the one case we have a 1019 mb. (30.09 ins.) ellipse travelling 5° to the westward, and in the other case a 1023 mb. (30.21 ins.) ellipse moving 5° to the eastward. This contrary travel in longitude may perhaps be accounted for by the fact that the sun is well south of the equator from December to February, making the African land relatively warmer than the ocean to the westward; whereas he is well to the north of the equator from July to September, making the African land in the same parallels as the anticyclone, relatively cooler than the ocean to the westward. The lower pressure over Africa in the first case would restrict the ellipse of higher pressure on the eastern side, and the higher pressure over Africa in the other case would extend it eastward. During these three months the 1019 mb. (30.09 ins.) isobar widens out, east and west, and overlaps the continents of Africa and America, and at the same time moves slightly to the northward. The isobar 1023 mb. (30.21 ins.) does not occupy a central position in regard to the isobar 1019 mb. (30.09 ins.), but is situated somewhat to the south thus steepening the gradient for westerly winds, between the 30th and 40th parallels. The region of south-east trade winds is more restricted, in the South Atlantic, than during any other period; but north of a straight line drawn from Port Nolloth on the African coast to Bahia in Brazil, the air current is still fairly constant from a south-easterly quarter, when it is not deflected by proximity to the land. Southward of this line to 35° S. the normal circulation round the central high pressure is to some extent maintained, but

MONTHLY CHARTS SHOWING THE RELATION OF THE ATLANTIC ANTICYCLONES TO THE REGION OF DOLDRUMS.  
(From the Barometer Manual for the use of Seamen, 8th Edition.)



The pressure represented by each isobar is in millibars, the equivalents of which are added.  
The extent and development of the region of Doldrums is shown by the area of shading.



MONTHLY CHARTS SHOWING THE RELATION OF THE ATLANTIC ANTICYCLONES TO THE REGION OF DOLDRUMS.  
(From the Barometer Manual for the use of Seamen, 8th Edition.)




Plate IV.

The pressure represented by each isobar is in millibars, the equivalents of which are added.  
The extent and development of the region of Doldrums is shown by the area of shading.

MONTHLY CHARTS SHOWING THE RELATION OF THE ATLANTIC ANTICYCLONES TO THE REGION OF DOLDRUMS.  
(From the Barometer Manual for the use of Seamen, 8th Edition.)





Plate V.

The pressure represented by each isobar is in millibars, the equivalents of which are added.  
The extent and development of the region of Doldrums is shown by the area of shading.





The pressure represented by each isobar is in millibars, the equivalents of which are added.  
The extent and development of the region of Doldrums is shown by the area of shading.

the winds are not constant; southwards of  $35^{\circ}$  S. the circulation becomes re-established and westerly winds predominate.

Over the eastern margin of the South Pacific a south-easterly air current largely predominates as far south as the 30th parallel, except where proximity to land induces a more southerly, or even a south-westerly, direction according to the trend of the coast-line.

*October and November.*—Over the South Atlantic north of the 30th parallel, as far west as the 10th meridian of west longitude, and north of the 15th parallel westward, the south-east trade wind is steady. West of the 10th meridian between the 15th and 30th parallels the tendency for northing in the prevailing winds due to the trend of the isobars becomes more marked the more westerly the position; south of the 30th parallel the winds have more westerly the more southerly the position. The occurrence of easterly gales from  $30^{\circ}$  W. to  $40^{\circ}$  W. between  $45^{\circ}$  S. and  $50^{\circ}$  S. is remarkable.

Over the eastern margin of the South Pacific northward of  $30^{\circ}$  S. the south-east trade wind is steady, and is fairly steady between  $30^{\circ}$  S. and  $35^{\circ}$  S. near the land, but its direction here, as to the northward, in the neighbourhood of the coast, is largely that of the coast-line.

#### AIR TEMPERATURE.

There is a general relation to be found between the direction of the prevailing wind and the average temperature of the air over the South Atlantic, on the eastern and western sides of the permanent area of high pressure.

A deviation of the isotherms from the east and west direction of the lines of latitude is found on the eastern and western sides of the high pressure area; the south-east trade wind produces a cooling effect upon the air on the eastern side, and the north-easterly wind a warming effect upon the air on the western side.

Throughout the year there is a marked tendency for air isotherms to extend along a line drawn from the Cape of Good Hope to the Island of Ascension, under the influence of the relatively cool air of the south-east trades which blow on the African side of the high pressure, and in association with the cool sea surface current setting north-westward. While, at the same time, the relatively warm north-east wind on the South American side of the anticyclone, which is in association with the warm sea surface current setting south-westward, exercises a similar influence upon the air isotherms in the vicinity of South America.

From December to February, inclusive, the south-east trade wind carries the eastern ends of the air isotherms considerably to the north-west of the average latitude for that temperature. The average maximum limit of effect for the three months is in about  $2^{\circ}$  N.  $18^{\circ}$  W., although the range during the three months is from  $5^{\circ}$  N.  $20^{\circ}$  W. to  $5^{\circ}$  S.  $15^{\circ}$  W.

Near the American coast the effect of the north-east wind is shown in the isotherms even as far to the southward as  $37^{\circ}$  S., ranging from  $40^{\circ}$  S. to  $35^{\circ}$  S. Consequently, on an average, the air isotherm of  $70^{\circ}$  F., for example, reaches the coast of Africa in  $17^{\circ}$  S., but dips to  $34^{\circ}$  S. on the coast of South America.



On the eastern margin of the South Pacific, as a general rule, the relatively cool southerly wind, corresponding with the prevailing cool sea surface current which sets to the northward, has an appreciable effect in lowering the temperature of the air near the west coast of South America. From December to February, inclusive, the cool southerly wind affects the direction of the air isotherms, forcing their easterly ends to the north-west, from  $40^{\circ}$  S. to  $5^{\circ}$  S., thus indicating a lower air temperature than is experienced along the same parallels of latitude farther to the westward.

Over the South Atlantic during the four months March to June, inclusive, the mean of extreme limits of the cooling effect of the south-east trades as indicated by the air isotherms, is  $3^{\circ}$  S.  $16^{\circ}$  W.; with a range from  $8^{\circ}$  S.  $15^{\circ}$  W. to  $5^{\circ}$  N.  $20^{\circ}$  W. The southern limit of influence of the north-east wind, on the western side of the ocean, gradually travels northward, month by month, from  $30^{\circ}$  S. to  $23^{\circ}$  S., and the average is found in about  $25^{\circ}$  S. The air isotherm of  $70^{\circ}$  F., on an average, terminates in  $16^{\circ}$  S., on the eastern side, but in  $29^{\circ}$  S. on the western side of the South Atlantic.

Over the eastern margin of the South Pacific during these months, the cooling effect is noticeable from  $40^{\circ}$  S. to the equator.

Over the South Atlantic throughout the three months July, August, September, the furthest extension of the cooling effect of the south-east trades is in about  $10^{\circ}$  N.  $20^{\circ}$  W.; and the warming effect of the north-east wind is at its southern limit in  $30^{\circ}$  S. The air isotherm of  $70^{\circ}$  F. terminates, on an average, in  $9^{\circ}$  S. on the coast of South Africa, and in  $25^{\circ}$  S. on the coast of South America.

Over the eastern margin of the South Pacific during this period the southern limit of effect due to the southerly wind is shown to have retreated  $5^{\circ}$  to the northward and is now in  $35^{\circ}$  S., while the cooling of the air can be traced to the equator.


Over the South Atlantic during October and November, the extreme limit of the effect of the south-east trades is in  $5^{\circ}$  N.  $20^{\circ}$  W., and the southern limit of that due to the north-east wind in  $30^{\circ}$  S. The isotherm of  $70^{\circ}$  F. terminates in  $15^{\circ}$  S. on the African coast, and in  $28^{\circ}$  S. on the American coast. At the same time, over the eastern margin of the South Pacific the cooling effect due to the southerly wind is evident from  $40^{\circ}$  S. to  $5^{\circ}$  S.

#### GALES.

Depressions of the South Atlantic which are attended with gale force appear to reach that ocean in two ways. They cross the continent of South America somewhere between  $25^{\circ}$  S. and Cape Horn, more frequently travelling over Patagonia; or they avoid the land altogether and round Cape Horn to the eastward, following the general drift of air and sea surface.

The percentage curves, *see* Plates VII. and VIII., seem to show that not infrequently a cyclonic system, instead of rounding Cape Horn or crossing the land, strikes northward or north-westward

#### GALE FREQUENCY.




The percentage of gales to the whole number of wind observations is indicated by the figure linked to each gale frequency line.



## GALE FREQUENCY.


JULY

AUGUST




SEPTEMBER

OCTOBER



NOVEMBER

DECEMBER



The percentage of gales to the whole number of wind observations is indicated by the figure linked to each gale frequency line.

up the South Pacific at a distance of  $5^{\circ}$  to  $10^{\circ}$  from the west coast of South America.

It seems probable that systems rounding Cape Horn drift north-east and east, their centres eventually pass south of the Cape of Good Hope, and continue their eastward movement in the Southern Ocean.

## Fog.

Attention may here be called to the effects of a warm moist wind over cold water, and of a cold wind over warm water.

A warm humid wind passing over water of a lower temperature is cooled, and the excess of vapour, after the point of saturation has been reached, may be condensed, and form mist or fog. In like manner a cold wind, blowing over water of a considerably higher temperature than its own, chills below the point of saturation the vapour as it rises, when it becomes visible. For the lower the temperature of the atmosphere the less aqueous vapour can it hold in an invisible state.

In the first case the fog, as a rule, does not extend more than a few hundred feet above the sea surface, and may occasionally be seen over from the masthead of a vessel; in the second case it rises to a comparatively high altitude, but does not always extend to the surface, so that objects may occasionally be seen at a distance by an observer descending to a position near sea-level. It must not be concluded that these are the only causes productive of fog at sea, but the regions of greatest fog frequency are regions in which these causes occur most conspicuously.

Over the South Atlantic, north of the 30th parallel, fog is rarely met with, except near the land on either side of the ocean. For over this area the wind circulates, for the most part, steadily round the central area of highest pressure, and the relation between air and sea surface temperature is fairly stable. Anywhere south of the 30th parallel, where a relatively warm moisture-laden wind passes over a relatively cold current, fog may be expected, and is found to become increasingly frequent the higher the latitude reached, to the southern limit of observation, see Plate IX. This is probably due to increase in gale frequency with latitude, the passage of cyclonic systems bringing in their train rapid fluctuations in air temperature, varying inequalities between the temperature of air and sea surface, together with the failing winds and calms which usually intervene between successive disturbances.

## THE DOLDRUMS.

Attention has already been called on p. 7 to the migration in the course of the year of the belt of calms and variable airs known as the doldrums of the equator. These are the regions between the two "trades," and they are localities of frequent rainfall. The weather of the doldrums is a subject of special interest on account of the part which it was supposed to take in the formation and maintenance of the trade-winds. A table giving a summary of observations on ships in that region is therefore given here.



THE STATE OF THE SKY AND THE WEATHER IN THE DOLDRUMS OF THE NORTH ATLANTIC IN THE SEVERAL MONTHS OF THE YEAR.

Cloud Amount ...	0-2	3-4		5-6		7-8		9-10		Sums.	Weather.			Sun's Mean Monthly Declination.		
		b		bc		cb		c			o		No. of obs.		d.h.r.p. No. of obs.	Wet weather per cent.
		No. of obs.	Per cent.	No. of obs.	Per cent.	No. of obs.	Per cent.	No. of obs.	Per cent.		No. of obs.	Per cent.				
Beaufort Notation...																
January	...	39	6.1	102	16.0	94	14.7	177	27.7	227	35.5	639	1,656	672	41	20° 18' S.
February	...	46	6.4	132	18.4	139	19.4	169	23.5	232	32.3	718	1,337	575	43	12° 44' S.
March	...	60	8.9	141	20.9	140	20.7	156	23.1	179	26.5	676	1,854	697	37	1° 58' S.
April	...	99	9.8	201	19.9	196	19.4	242	23.9	275	27.1	1,013	1,605	746	46	9° 27' N.
May	...	58	5.6	173	16.5	219	20.8	257	24.4	345	32.7	1,052	2,060	927	45	18° 22' N.
June	...	63	6.2	163	15.9	204	19.9	247	24.1	348	33.9	1,025	1,842	977	53	22° 36' N.
July	...	87	8.7	205	19.3	210	19.7	266	25.0	295	27.7	1,063	2,116	860	41	20° 48' N.
August	...	114	10.0	285	25.0	272	23.9	238	20.9	228	20.0	1,137	1,744	596	34	13° 32' N.
September	...	213	14.0	406	26.8	299	19.7	310	20.4	287	19.0	1,515	2,403	772	32	3° 2' N.
October	...	150	13.5	264	23.7	237	21.3	237	21.3	226	20.3	1,114	2,718	1,090	40	8° 26' S.
November	...	49	6.1	149	18.6	153	19.1	169	21.1	280	35.0	800	2,008	1,043	50	17° 54' S.
December	...	36	7.4	84	17.3	100	20.6	126	26.0	139	28.7	485	1,161	447	38	22° 26' S.
Year	...	1,014	9.0	2,305	20.5	2,263	20.1	2,594	23.1	3,061	27.3	11,237	22,504	9,402	42	


It will be noticed that the sky is practically free from cloud for more than ten per cent. of the observations in September, October and November; on the other hand May is the month which is least favoured with the weather denoted by *b*.

The number of observations of overcast sky varies irregularly over the year from 35.5 per cent. in January, 35 per cent. in November to 19 per cent. in September and 20 per cent. in August.

Including all forms of rainy weather *d*, *h*, *r*, *p*, 53 per cent. of the observations are identified with rain in June and 32 per cent. in August; the average for all the months is 42 per cent.

## DISTRIBUTION AND FREQUENCY OF FOG.


OCTOBER TO FEBRUARY.



The upper figures show the number of wind observations, the lower the percentage of those which are associated with Fog.

## DISTRIBUTION AND FREQUENCY OF FOG.

MARCH TO SEPTEMBER.



The upper figures show the number of wind observations, the lower the percentage of those which are associated with Fog.



## INDEX.

	PAGE
Air circulation, South Atlantic ...	5
Air temperature ...	9
Anticyclone, South Atlantic ...	5
"    Winds of ...	5-6
Argentina, Daily Weather Report of ...	6
Barometer, Wind and ...	5
Barometric pressure, effect of littoral on ...	7
"    "    and passage of depressions ...	6
Brazil coast, Low pressure over ...	6
British Association ...	3
Cape Horn, Wind systems rounding ...	10
Cause of fog, A ...	11
Chart of South Atlantic wind ...	5
Circulation of air over South Atlantic ...	5
Coasts and trend of isobars ...	7
Cooling effect of southerly wind ...	9, 10
Council, Meteorological... ..	3
Cyclones and temperature fluctuations ...	11
Daily Weather Charts, Argentina ...	6
Depressions, Eastward movement of ...	6
Deviation of isotherms ...	9
Distribution of wind, pressure, and temperature ...	7-9
Doldrums, Mean latitude of ...	7
"    Rainfall in ...	12
"    State of sky in ...	12
Eastward moving depressions ...	6
Elliptical isobars... ..	7
Fog ...	11
"    Cause of ...	11
"    distribution... ..	11
"    frequency ...	11
Frequency of gales ...	7, 10
Gale frequency ...	7, 10
Gales near Brazilian coast ...	6
"    of South Atlantic ...	10
Gradient and wind ...	7
Hydrographer of the Navy ...	3
Influence of littoral on wind direction ...	7, 9
Isobars, Elliptical ...	7
Isotherms, Deviation of ...	9
"    Influence of the winds on... ..	9
"    Trend of ...	9
Littoral, Wind follows ...	7, 9
Low pressure, Coast of Brazil ...	6
"    South Africa ...	5
"    systems travelling eastward ...	6
Major axis of South Atlantic high pressure ...	7
Maximum gale force of South Atlantic ...	6
Mean latitude of doldrum belt... ..	7
Meteorological Council ...	3
Pressure distribution, Barometric ...	7-9
"    Major axis of high ...	7
Rainfall in doldrums ...	12
Rear-Admiral Sir W. J. L. Wharton ...	3
Relation between temperature, pressure and air circulation ...	5
"    wind and gradient ...	7
River Plate, low pressure ...	6



	PAGE
South Africa, low pressure ... ..	5
South Atlantic, air circulation ... ..	5
" " " temperature ... ..	9
" " gales ... ..	10
" " maximum gale force ... ..	6
" " wind chart ... ..	5
" " " systems ... ..	5
South-east trade winds ... ..	7-9
Southerly wind, Cooling effect of ... ..	9, 10
South Pacific, air temperature ... ..	10
" " anticyclone ... ..	7
Systems rounding Cape Horn ... ..	10
Temperature fluctuations and cyclones ... ..	11
Temperature of the air ... ..	9
Trade winds ... ..	8
Travelling low pressure systems ... ..	6
Travel of high pressure systems ... ..	8
Trend of isobars with coast line ... ..	7
Wind and barometer ... ..	5
Wind and gradient ... ..	7
" " temperature ... ..	9
" chart of South Atlantic... ..	5
" distribution ... ..	6
" follows littoral ... ..	7, 9
" maximum force of South Atlantic ... ..	6
Wharton, Rear-Admiral Sir W. J. L....	3
Winds of anticyclone ... ..	5-6
Wind systems of South Atlantic ... ..	5



W.  
Wind sy